

Eigenvalue estimates for submanifolds with locally bounded mean curvature

G. Pacelli Bessa

J. Fábio Montenegro

February 1, 2008

Abstract

We give lower bounds estimates for the first Dirichlet eigenvalues for domains Ω in submanifolds $M \subset N$ with locally bounded mean curvature. These lower bounds depend on the local injectivity radius, local upper bound for sectional curvature of N and local bound for the mean curvature of M . For submanifolds with bounded mean curvature of Hadamard manifolds these lower bounds depends only on the dimension and the bound on mean curvature.

1 Introduction

Let M be a Riemannian manifold and $\Omega \subset M$ a connected domain with compact closure and nonempty boundary $\partial\Omega$. The first eigenvalue of Ω denoted by $\lambda_1(\Omega)$ is given by

$$\lambda_1(\Omega) = \inf \left\{ \frac{\int_{\Omega} |\nabla f|^2}{\int_{\Omega} f^2}, f \in L^2_{1,0}(\Omega) \setminus \{0\} \right\}, \quad (1)$$

where $L^2_{1,0}(\Omega)$ is the completion of $C_0^\infty(\Omega)$ with respect to the norm

$$\|\varphi\|^2 = \int_{\Omega} \varphi^2 + \int_{\Omega} |\nabla \varphi|^2. \quad (2)$$

Observe that in this variational formulation, the boundary $\partial\Omega$ needs not to be smooth. It is clear that if $\Omega_1 \subset \Omega_2$ are bounded domains, then $\lambda_1(\Omega_1) \geq \lambda_1(\Omega_2) \geq 0$. From this considerations one may define the first eigenvalue of a complete non-compact Riemannian manifold M as follows.

Definition 1.1 *The first eigenvalue of a complete non-compact Riemannian manifold M is*

$$\lambda_1(M) = \lim_{r \rightarrow \infty} \lambda_1(B_M(p, r)), \quad (3)$$

where $B_M(p, r)$ is the geodesic ball of radius r centered at p .

This limit (3) exists and it is independent on the center p . Since $\lambda_1(M)$ can be zero, it is natural to ask under what conditions a complete non-compact Riemannian manifold M has positive first eigenvalue. Toward this question, McKean proved the following theorem.

Theorem 1.2 (Mckean, [6]) *Let M be an m -dimensional, complete non-compact, simply connected Riemannian manifold with sectional curvature $K_M \leq -a^2 < 0$, then*

$$\lambda_1(M) \geq \frac{(m-1)^2 a^2}{4} \quad (4)$$

Leung-Fu Cheung and Pui-Fai Leung in [3] considered non-compact m -dimensional submanifolds M of the hyperbolic space $\mathbb{H}^n$ with bounded mean curvature H and showed that if $|H| \leq c$ for some constant $c < m-1$, then $\lambda_1(M) \geq (m-1-c)^2/4$. They considered the distance function r (distance to a point) in the hyperbolic space $\mathbb{H}^n(-1)$ and showed that when r is restricted to a non-compact submanifold M with mean curvature $|H| \leq c < n-1$ then $\Delta r \geq n-1-c$, where Δ is the laplace operator in the induced metric on M . After that, using cleverly Young inequality concluded that $\lambda_1(M) \geq (n-1-c)^2/4$.

In this paper we will consider similar questions regarding eigenvalues estimates. Let $M \subset_\varphi N$ be an immersed submanifold with locally bounded mean curvature in a Riemannian manifold N (see section 4 for a definition). Let $p \in N$ and consider connected components $\Omega \subset \varphi^{-1}(B_N(p, r))$. We give lower bounds estimates for the first Dirichlet eigenvalue $\lambda_1(\Omega)$. The radius of the ball and the lower bounds depend on the sectional curvatures of the N in $B_N(p, r)$, on the injectivity radius $inj(p)$ of N at p and the mean curvature of Ω (see Theorem (4.2)). When N is complete, simply connected and has sectional curvature bounded from above by a negative constant the lower bounds for $\lambda_1(\Omega) \geq c > 0$, where c is independent on r . Thus, we can conclude that $\lambda_1(M) > c$ (see Corollary 4.4). This extends Cheun-Leung result to submanifolds with bounded mean curvature of Hadamard manifolds.

2 Main Lemma (Lemma 2.1)

As mentioned before, Cheung-Leung showed that the distance function r when restricted to a non-compact m -dimensional submanifold $M \subset \mathbb{H}^n(-1)$ with mean curvature $|H| < m-1$, the laplacian of r in the induced metric on M is bounded from below, $\Delta r \geq n-1-\sup |H|$. Using Young inequality they showed that $\lambda_1(M) \geq (n-1-\sup |H|)^2/4$. Their idea (and proof) applies immediately to vector fields X on domains Ω of manifolds provided that $\inf \operatorname{div} X > 0$ and $\sup |X| < \infty$ on Ω . This is the content of the following lemma.

Lemma 2.1 *Let $\Omega \subset M$ be a domain with compact closure ($\partial\Omega \neq \emptyset$) in a Riemannian manifold M and $X : M \rightarrow TM$ be a smooth nonzero vector field with $\sup_\Omega |X| < \infty$ and $\inf_\Omega \operatorname{div} X > 0$, then*

$$\lambda_1(\Omega) \geq \left[\frac{\inf_\Omega \operatorname{div} X}{2 \sup_\Omega |X|} \right]^2 > 0. \quad (5)$$

Proof: Let $f \in C_0^\infty(\Omega)$. The vector field $f^2 X$ has compact support on Ω . Computing $\operatorname{div}(f^2 X)$ we have

$$\begin{aligned} \operatorname{div}(f^2 X) &= \langle \operatorname{grad} f^2, X \rangle + f^2 \operatorname{div}(X) \\ &\geq -|\operatorname{grad} f^2| \cdot |X| + \inf \operatorname{div} X \cdot f^2 \\ &\geq -2 \cdot \sup |X| \cdot |f| |\operatorname{grad} f| + \inf \operatorname{div} X \cdot f^2 \end{aligned} \quad (6)$$

Using the inequality

$$-2|f| \cdot |\operatorname{grad} f| \geq -\epsilon \cdot |f|^2 - 1/\epsilon \cdot |\operatorname{grad} f|^2$$

for all $\epsilon > 0$, we have from (6) that

$$\operatorname{div}(f^2 X) \geq \sup |X| \cdot (-\epsilon \cdot |f|^2 - \frac{1}{\epsilon} \cdot |\operatorname{grad} f|^2) + \inf \operatorname{div} X \cdot f^2 \quad (7)$$

Integrating (7) over Ω we have that

$$0 = \int_{\Omega} \operatorname{div}(f^2 X) \geq \sup |X| \cdot \int_{\Omega} (-\epsilon |f|^2 - \frac{1}{\epsilon} |\operatorname{grad} f|^2) + \inf \operatorname{div} X \cdot \int_{\Omega} f^2,$$

therefore

$$\int_{\Omega} |\operatorname{grad} f|^2 \geq \frac{\epsilon}{\sup |X|} (\inf \operatorname{div} X - \sup |X| \cdot \epsilon) \int_{\Omega} f^2.$$

Choosing $\epsilon = (\inf \operatorname{div} X)/(2 \sup |X|)$ we have that

$$\int_{\Omega} |\operatorname{grad} f|^2 \geq \left[\frac{\inf \operatorname{div} X}{2 \sup |X|} \right]^2 \int_{\Omega} f^2. \quad (8)$$

By the variational formulation of the first eigenvalue of Ω the inequality (8) implies (5).

Corollary 2.2 *There are no smooth bounded vector fields $X : M \rightarrow TM$ with $\inf_M \operatorname{div} X > 0$ on complete non-compact manifolds with $\lambda_1(M) = 0$. In particular there is no such vector fields on $\mathbb{R}^n$.*

3 Basic Formulas

In order to apply Lemma (2.1) we need to find suitable vector fields $X \in TN$ such that the divergent of the tangential component $X^t \in TM \subset TN$ has positive infimum. The natural vector field is the gradient of the distance function that depending on the geometry of M and N has divergent strictly positive. To see this we shall set up some basic formulas relating the divergent of the gradient of the distance function, its lower bound, the mean curvature and upperbounds of the sectional curvatures of N .

Let $\varphi : M \hookrightarrow N$ be an isometric immersion, where M and N are complete Riemannian manifolds. Consider a smooth function $g : N \rightarrow \mathbb{R}$ and the composition $f = g \circ \varphi : M \rightarrow \mathbb{R}$. Identifying X with $d\varphi(X)$ we have at $p \in M$ and for every $X \in T_p M$ that

$$\langle \operatorname{grad} f, X \rangle = df(X) = dg(X) = \langle \operatorname{grad} g, X \rangle,$$

therefore

$$\operatorname{grad} g = \operatorname{grad} f + (\operatorname{grad} g)^\perp, \quad (9)$$

where $(\operatorname{grad} g)^\perp$ is perpendicular to $T_p M$. Let ∇ and $\bar{\nabla}$ be the Riemannian connections on M and N respectively, $\alpha(p)(X, Y)$ and $Hess f(p)(X, Y)$ be respectively the second fundamental form of the immersion φ and the Hessian of f at $p \in M$, $X, Y \in T_p M$. Using the Gauss equation we have that

$$Hess f(p)(X, Y) = Hess g(\varphi(p))(X, Y) + \langle \operatorname{grad} g, \alpha(X, Y) \rangle_{\varphi(p)}. \quad (10)$$

Now let $\{e_1, \dots, e_m\}$ be an orthonormal basis for $T_p M$. Then taking the trace in (10) we have that

$$\begin{aligned}\Delta f(p) &= \sum_{i=1}^m \text{Hess } f(p)(e_i, e_i) \\ &= \sum_{i=1}^m \text{Hess } g(\varphi(p))(e_i, e_i) + \langle \text{grad } g, \sum_{i=1}^m \alpha(e_i, e_i) \rangle.\end{aligned}\quad (11)$$

We should mention that the formulas (10) and (11) are well known in the literature, see [3], [4], [2]. Another important tool is the Hessian Comparison Theorem.

Theorem 3.1 (Hessian Comparison Theorem) *Let M be a complete Riemannian manifold and $x_0, x_1 \in M$. Let $\gamma : [0, \rho(x_1)] \rightarrow M$ be a minimizing geodesic joining x_0 and x_1 where $\rho(x)$ is the distance function $\text{dist}_M(x_0, x)$. Let K be the sectional curvatures of M and $\mu_i(\rho)$, $i = 0, 1$, be these functions defined below.*

$$\mu_0(\rho) = \begin{cases} k_0 \cdot \coth(k_0 \cdot \rho(x)), & \text{if } \inf_{\gamma} K = -k_0^2 \\ \frac{1}{\rho(x)}, & \text{if } \inf_{\gamma} K = 0 \\ k_0 \cdot \cot(k_0 \cdot \rho(x)), & \text{if } \inf_{\gamma} K = k_0^2 \text{ and } \rho < \pi/2k_0. \end{cases} \quad (12)$$

And

$$\mu_1(\rho) = \begin{cases} k_1 \cdot \coth(k_1 \cdot \rho(x)), & \text{if } \sup_{\gamma} K = -k_1^2 \\ \frac{1}{\rho(x)}, & \text{if } \sup_{\gamma} K = 0 \\ k_1 \cdot \cot(k_1 \cdot \rho(x)), & \text{if } \sup_{\gamma} K = k_1^2 \text{ and } \rho < \pi/2k_1. \end{cases} \quad (13)$$

Then the Hessian of ρ and ρ^2 satisfies

$$\mu_0(\rho(x)) \cdot |X|^2 \geq \text{Hess } \rho(x)(X, X) \geq \mu_1(\rho(x)) \cdot |X|^2 \quad (14)$$

$$2\rho(x) \cdot \mu_0(\rho(x)) \cdot |X|^2 \geq \text{Hess } \rho^2(x)(X, X) \geq 2\rho(x) \cdot \mu_1(\rho(x)) \cdot |X|^2.$$

Where X is any vector in $T_x M$ perpendicular to $\gamma'(\rho(x))$. Taking the trace in (14) we have

$$(n-1)\mu_0(\rho(x)) \geq \Delta \rho(x) \geq (n-1)\mu_1(\rho(x)) \quad (15)$$

$$2(n-1)\rho(x)\mu_0(\rho(x)) \geq \Delta \rho^2(x) \geq 2(n-1)\rho(x)\mu_1(\rho(x)).$$

Here $n = \dim M$.

4 Applications of the Main Lemma

Let N be a complete Riemannian manifold and let $\kappa(p, r) = \sup\{K_N(x); x \in B_N(p, r)\}$, where $K_N(x)$ are the sectional curvatures of N at x and $B_N(p, r)$ is the geodesic ball with center p and radius r . As an application of the lemma (2.1) we are going to give Dirichlet eigenvalue lower bound estimate for domains on immersed submanifolds with locally bounded mean curvature.

Definition 4.1 *An isometric immersion $\varphi : M \hookrightarrow N$ has locally bounded mean curvature H if for any $p \in N$ and $r > 0$ this number $h(p, r) = \sup\{|H(x)|; x \in \varphi(M) \cap B_N(p, r)\}$ is finite.*

Cheng in ([1]) gave lower bounds for the first Dirichlet eigenvalue on balls $B_N(p, r)$, $r < \text{inj}(p)$. These bounds are the first Dirichlet eigenvalue on balls of the same radius on space forms with constant curvature $\sup_{B_N(p, r)} K_N$. The following theorem can be viewed as a version of Cheng's eigenvalue estimates for balls in submanifolds.

Theorem 4.2 *Let $\varphi : M \hookrightarrow N$ be an isometric immersion with locally bounded mean curvature H where M is an m -dimensional complete non-compact Riemannian manifold. Choose any point $p \in N \setminus \varphi(M)$ and r as follows:*

Case I. The injectivity radius of N at p is finite, $\text{inj}(p) < \infty$.

1. *If $\kappa(p, \text{inj}(p)) = k^2$, choose $r < \min\{\text{inj}(p), \pi/2k, \cot^{-1}[h(p, \text{inj}(p))/(m-1)k]/k\}$.*

2. *If $\kappa(p, \text{inj}(p)) = 0$, choose $r < \min\{\text{inj}(p), (m-1)/h(p, \text{inj}(p))\}$.*

3. *If $\kappa(p, \text{inj}(p)) = -k^2$ and $h(p, \text{inj}(p)) < m-1$, then choose $r < \text{inj}(p)$.*

If $h(p, \text{inj}(p)) \geq m-1$ then choose $r < \min\{\text{inj}(p), \coth^{-1}[h(p, \text{inj}(p))/(m-1)k]/k\}$

Case II. The injectivity radius of N at p is infinity, $\text{inj}(p) = \infty$.

4. *If $\lim_{r \rightarrow \infty} \kappa(p, r) = \infty$, choose $r = \max_{s>0} \min\{\frac{\pi}{2\sqrt{\kappa(p, s)}}, \cot^{-1}[\frac{h(p, s)}{(m-1)k(p, 2)}]/k(p, s)\}$.*

5. *If $\lim_{r \rightarrow \infty} \kappa(p, r) = 0, -k^2$, choose r as in 2. and 3. respectively.*

Let Ω be any connected component of $\varphi^{-1}(\overline{B_N(p, r)})$, r chosen above. Then the first eigenvalue $\lambda_1(\Omega)$ for the Dirichlet problem on Ω has the following lower estimates.

- *r as in 1. and 4.: $\lambda_1(\Omega) \geq [(m-1)k \cot(kr) - h(p, r)]^2/4$, where in 4. we substitute k by $\sqrt{k(p, r)}$.*

- *r as in 2.: $\lambda_1(\Omega) \geq [(m-1)/r - h(p, r)]^2/4$*

- *r as in 3.: If $h(p, \text{inj}(p)) < m-1$ then $\lambda_1(\Omega) \geq [(m-1) - h(p, r)]^2/4$.*

If $h(p, \text{inj}(p)) \geq m-1$ then $\lambda_1(\Omega) \geq [(m-1)k \coth(kr) - h(p, r)]^2/4$.

Proof: Let $\rho(x) = \text{dist}_N(p, x)$ be the distance function on N and $f = \rho \circ \varphi : M \rightarrow \mathbb{R}$. f is smooth on $\varphi^{-1}(B_N(p, \text{inj}(p)))$. Consider a connected component Ω of $\varphi^{-1}(\overline{B_N(p, r)})$ and in this component let $X = \text{grad } f$. Now by (11), $\text{div } X(x) = \sum_{i=1}^{m-1} \text{Hess } \rho(\varphi(x))(e_i, e_i) + \langle \text{grad } \rho, H \rangle(\varphi(x))$, $\{e_1, \dots, e_m\}$ base orthonormal of $T_x M$. Applying the Hessian Comparison Theorem we have that

- *$\text{div } X \geq (m-1)k \cot(kr) - h(p, r) > 0$, if $\kappa(p, r) = k^2$.*

- $\operatorname{div} X \geq (m-1)/r - h(p, r) > 0$, if $\kappa(p, r) = 0$.
- $\operatorname{div} X \geq (m-1)k \coth(kr) - h(p, r) > 0$, if $\kappa(p, r) = -k^2$.

Since $|X| \leq 1$ the estimative follows from Lemma (2.1).

Remark 4.3 *It was proved by Jorge and Xavier in [5] (Lemma 1) that if $\varphi : M \rightarrow N$ is an isometric immersion with locally bounded second fundamental form α and r chosen as in Theorem (4.2) where $h(p, r)$ is substituted by $\beta(p, r) = \{|\alpha(x)|; x \in \varphi(M) \cap B_N(p, r)\}$, then $\varphi^{-1}(\overline{B_N(p, r)})$ is a countable collection of disjoint compact topological m -disks Ω_j and the function $f = \rho \circ \varphi$ has only one critical point $q_j \in \Omega_j$ and its a minimum. Moreover, $\operatorname{diam}_M(\Omega_j)$ is bounded above by $C(r, \rho(\varphi(q_j)))$. It also shwon that Ω_j contains a geodesic ball centered at q_j with radius $r_j \geq c(r, \rho(\varphi(q_j)), \inf_{B_N(p, r)} K_N)$.*

Corollary 4.4 *Let $\varphi : M \hookrightarrow N$ be an isometric immersion with bounded mean curvature H , where M is an m -dimensional complete non-compact Riemannian manifold M and N is an n -dimensional complete simply connected Riemannian manifold N is with sectional curvature K_N satisfying*

$$K_N \leq -a^2 < 0, \text{ for a positive constant } a > 0.$$

If $|H| \leq \beta < (m-1)a$, then

$$\lambda_1(M) \geq \frac{[(m-1)a - \beta]^2}{4} > 0. \quad (16)$$

In particular, there is entire Green's functions on M .

Proof: As in Theorem (4.2), consider $\rho = \operatorname{dist}_N(p, \cdot)$, $p \in N \setminus \varphi(M)$, $f = \rho \circ \varphi$ and $X = \operatorname{grad} f$. Since N is simply connected and is negatively curved, every point of N is a pole, hence $\operatorname{inj}(p) = \infty$. The estimative for the case $\operatorname{inj}(p) = \infty$ and $h(p, r) < m-1$ for all $r > 0$ in Theorem (4.2) does depend on r , i.e. $\lambda_1(\Omega) \geq (m-1-\beta)^2/4$. Since any domain $\Omega \subset M$ is immersed inside a ball $B_N(p, r)$ for sufficient large r we have that $\lim \lambda_1(\Omega_j) \geq (m-1-\beta)^2/4$ for any exhaustion $\{\Omega_j \subset \Omega_{j+1}\}$ of M . The existence of entire Green's functions on M is a direct consequence of Theorem A.3 from [7], page 84.

References

- [1] Cheng, S. Y., *Eigenfunctions and eigenvalues of the Laplacian*. Am. Math. Soc Proc. Symp. Pure Math. 27, part II, 185-193 (1975).
- [2] Choe, J. and Gulliver, R., *Isoperimetric inequalities on minimal submanifolds of space forms*. Manuscripta Math. 77 (1992) 169-189.
- [3] Cheung, Leung-Fu and Leung, Pui-Fai, *Eigenvalue estimates for submanifolds with bounded mean curvature in the hyperbolic space*. Math. Z 236, 525-530 (2001)
- [4] Jorge, L. and Koutrofotis, D., *An estimate for the curvature of bounded submanifolds*. American Journal of Mathematics, vol. 103, n^o4 711-725 (1980)

- [5] Jorge, L. and Xavier, F., *An inequality between the exterior diameter and the mean curvature of bounded immersions*. Math. Z. 178, 77-82(1981).
- [6] McKean, H. P., *An upper bound for the spectrum of Δ on a manifold of negative curvature*. J. Differ. Geom. 4, 359-366(1970).
- [7] Schoen, R. and Yau, S. T., *Lectures on Differential Geometry*. Conference Proceedings and Lecture Notes in Geometry and Topology, vol.1